

Finding Order in Metric Structures (Part 2)

Aaron Anderson (joint with Diego Bejarano)

UPenn

May 4, 2026



Metric Structures

Definition

A *metric language* is just like a regular first-order language, consisting of functions and relations.

Definition

A *metric structure* consists of:

- A complete metric space of diameter ≤ 1
- For each n -ary function symbol, a uniformly continuous function $M^n \rightarrow M$
- For each n -ary relation symbol, a uniformly continuous function $M^n \rightarrow [0, 1]$

Formulas

Definition

An *atomic formula* is defined as usual, except instead of $=$, the basic relation is $d(x, y)$.

Definition

A *formula* is

- An atomic formula
- $u(\phi_1, \dots, \phi_n)$ where ϕ_i s are formulas and $u : [0, 1]^n \rightarrow [0, 1]$ is continuous
- $\sup_x \phi$ or $\inf_x \phi$

Making Linear Orders Metric

- Most famous examples of metric structures are stable or not NIP - how do we put an order on one?

Making Linear Orders Metric

- Most famous examples of metric structures are stable or not NIP - how do we put an order on one?
- Itai Ben Yaacov described Ordered Real Closed Metric Valued Fields, but making the metric space bounded complicated things.

Making Linear Orders Metric

- Most famous examples of metric structures are stable or not NIP - how do we put an order on one?
- Itai Ben Yaacov described Ordered Real Closed Metric Valued Fields, but making the metric space bounded complicated things.
- Diego Bejarano and I have a simpler approach.

Metric Linear Orders

- Call M a *metric linear order* if
 - M has a complete metric of diameter ≤ 1
 - M has a linear order
 - open balls are order-convex.
- M is a metric structure in the language $\{r\}$, with

$$r(x, y) = \begin{cases} 0 & x \leq y \\ d(x, y) & y \leq x \end{cases}$$

- Think of $r(x, y)$ as “the amount x is greater than y ,” or

$$r(x, y) = d(x, (\infty, y]).$$

Axiomatizing Metric Linear Orders

Definition

A *theory* is a set of *conditions* $\phi(x) = 0$.

Axiomatizing Metric Linear Orders

Definition

A *theory* is a set of *conditions* $\phi(x) = 0$.

Theorem (A., Bejarano)

Metric linear orders are axiomatized in $\{r\}$ by

$$\sup_{x,y} |(r(x,y) + r(y,x)) - d(x,y)| = 0 \quad (\text{antisymmetry})$$

$$\sup_{x,y} \min\{r(x,y), r(y,x)\} = 0 \quad (\text{linearity})$$

$$\sup_{x,y,z} r(x,z) - (r(x,y) + r(y,z)) = 0 \quad (\text{transitivity})$$

Ultrametric Dense Linear Orders

- DLO is a useful completion of the theory of linear orders

Ultrametric Dense Linear Orders

- DLO is a useful completion of the theory of linear orders
- We seek analogous completions of MLO

Ultrametric Dense Linear Orders

- DLO is a useful completion of the theory of linear orders
- We seek analogous completions of MLO
- To start, assume the metric is an **ultrametric**.

Axiomatizing Ultrametric Dense Linear Orders

Definition

Let UDLO be the theory of *ultrametric-dense linear orders*, consisting of MLO with the following axioms:

- $d(x, z) \leq \max(d(x, y), d(y, z))$
- For any rational $p \in \mathbb{Q} \cap [0, 1]$, $\sup_x \inf_y |r(x, y) - p| = 0$
- For any rational $p \in \mathbb{Q} \cap [0, 1]$, $\sup_x \inf_y |r(y, x) - p| = 0$.

Basically, the distances from x to $y > x$ are dense in $[0, 1]$.

Stable Diversion: Dense Ultrametrics

Call an ultrametric space *dense* if the set of distances to any point is dense in $[0, 1]$.

Fact (Conant)

The theory of dense ultrametrics is complete and has QE, but is not \aleph_0 -categorical.

Stable Diversion: Dense Ultrametrics

Call an ultrametric space *dense* if the set of distances to any point is dense in $[0, 1]$.

Fact (Conant)

The theory of dense ultrametrics is complete and has QE, but is not \aleph_0 -categorical.

(A separable ultrametric space has countably many distances!)

Stable Diversion: Dense Ultrametrics

Call an ultrametric space *dense* if the set of distances to any point is dense in $[0, 1]$.

Fact (Conant)

The theory of dense ultrametrics is complete and has QE, but is not \aleph_0 -categorical.

(A separable ultrametric space has countably many distances!)

Lemma (A., Bejarano)

The theory of dense ultrametrics is the model companion of the theory of ultrametrics, and is approximately \aleph_0 -categorical.

Ultrametric Dense Linear Orders

Theorem (A., Bejarano)

- UDLO *is complete*
- UDLO *eliminates quantifiers*
- UDLO *is the model companion of the theory of ultrametric linear orders*
- UDLO *is approximately \aleph_0 -categorical*

Ultrametric Dense Linear Orders

Theorem (A., Bejarano)

- UDLO *is complete*
- UDLO *eliminates quantifiers*
- UDLO *is the model companion of the theory of ultrametric linear orders*
- UDLO *is approximately \aleph_0 -categorical*

Theorem (A., Bejarano)

In a model of UDLO,

- *the metric and order topologies agree*
- *dcl = acl = topological closure.*

A Separable UDLO

- Let $S \subseteq (0, 1]$ be countable, dense.

A Separable UDLO

- Let $S \subseteq (0, 1]$ be countable, dense.
- Let $U_S = \{f : S \rightarrow \mathbb{Q} : \forall r > 0, \{s > r : f(s) \neq 0\} \text{ is finite.}\}$.

A Separable UDLO

- Let $S \subseteq (0, 1]$ be countable, dense.
- Let $U_S = \{f : S \rightarrow \mathbb{Q} : \forall r > 0, \{s > r : f(s) \neq 0\} \text{ is finite.}\}$.
- Let $d(f, g) = \max\{x : f(x) \neq g(x)\}$.

A Separable UDLO

- Let $S \subseteq (0, 1]$ be countable, dense.
- Let $U_S = \{f : S \rightarrow \mathbb{Q} : \forall r > 0, \{s > r : f(s) \neq 0\} \text{ is finite.}\}$.
- Let $d(f, g) = \max\{x : f(x) \neq g(x)\}$.
- Let $f(x) < g(x)$ when $f(d(f, g)) < g(d(f, g))$.

A Separable UDLO

- Let $S \subseteq (0, 1]$ be countable, dense.
- Let $U_S = \{f : S \rightarrow \mathbb{Q} : \forall r > 0, \{s > r : f(s) \neq 0\} \text{ is finite.}\}$.
- Let $d(f, g) = \max\{x : f(x) \neq g(x)\}$.
- Let $f(x) < g(x)$ when $f(d(f, g)) < g(d(f, g))$.

Lemma (A., Bejarano)

$U_S \models \text{UDLO}$.

A Separable UDLO

- Let $S \subseteq (0, 1]$ be countable, dense.
- Let $U_S = \{f : S \rightarrow \mathbb{Q} : \forall r > 0, \{s > r : f(s) \neq 0\} \text{ is finite.}\}$.
- Let $d(f, g) = \max\{x : f(x) \neq g(x)\}$.
- Let $f(x) < g(x)$ when $f(d(f, g)) < g(d(f, g))$.

Lemma (A., Bejarano)

$U_S \models \text{UDLO}$.

Theorem (Van Thé)

U_S has extremely amenable automorphism group, following from Fraïssé theory in the **discrete-logic** language of ordered S -valued metric spaces.

o-Minimality in Discrete Logic

Fact

If M expands a linear order, TFAE:

- every formula $\phi(x)$ in one variable is *qf-definable* in $\{<\}$
 - every formula $\phi(x)$ in one variable is a finite union of intervals.
-
- If these happen, M is *o-minimal*.

o-Minimality in Discrete Logic

Fact

If M expands a linear order, TFAE:

- every formula $\phi(x)$ in one variable is *qf-definable* in $\{<\}$
 - every formula $\phi(x)$ in one variable is a finite union of intervals.
-
- If these happen, M is *o-minimal*.
 - How do we describe these properties for MLOs?

Metric o-Minimality

Theorem (A., Bejarano)

If M expands a metric linear order, TFAE:

- *every formula $\phi(x)$ in one variable is qf-definable in $\{r\}$*
- *every formula $\phi(x)$ in one variable is regulated (a uniform limit of step functions).*

Metric o -Minimality

Theorem (A., Bejarano)

If M expands a metric linear order, TFAE:

- every formula $\phi(x)$ in one variable is qf -definable in $\{r\}$
- every formula $\phi(x)$ in one variable is regulated (a uniform limit of step functions).

Definition

If M expands a metric linear order, call M o -minimal if every formula $\phi(x)$ in one variable satisfies these equivalent properties.

Metric \mathfrak{o} -Minimality

Theorem (A., Bejarano)

If M expands a metric linear order, TFAE:

- *every formula $\phi(x)$ in one variable is qf -definable in $\{r\}$*
- *every formula $\phi(x)$ in one variable is regulated (a uniform limit of step functions).*

Definition

If M expands a metric linear order, call M *\mathfrak{o} -minimal* if every formula $\phi(x)$ in one variable satisfies these equivalent properties.

By QE, a model of UDLO is \mathfrak{o} -minimal.

Regulated Functions

Regulated functions $[a, b] \rightarrow \mathbb{R}$ were defined by Bourbaki.

Lemma (A. Bejarano)

If M is a linear order, $f : M \rightarrow [0, 1]$, TFAE:

- ① f is a uniform limit of step functions
- ② For any $a < b$, M can be partitioned into finitely many intervals on which either $f(x) > a$ or $f(x) < b$.

If $M \models \text{UDLO}$, and $f : M \rightarrow M$ is definable ($d(f(x), y)$ is a formula), then f is continuous and satisfies (2).

Metric Valued Fields

Definition

A *metric valued field* is a field K with an absolute value $|\cdot| : K \rightarrow \mathbb{R}^{\geq 0}$ satisfying valuation axioms:

- $|x| = 0 \iff x = 0$
- $|xy| = |x||y|$
- $|x + y| \leq \max(|x|, |y|)$

which gives rise to an ultrametric $d(x, y) = |x - y|$.

Real Closed Metric Valued Fields

Definition

If K is a metric valued field equipped with an order, $K \equiv \mathbb{R}$ as ordered fields in classical logic, and $|\cdot|$ takes values outside $\{0, 1\}$, call K a *real closed metric valued field*.

These are almost metric structures, but the metric is unbounded, a problem for continuous logic.

Metric Valued Fields, à Rideau-Kikuchi, Scanlon, Simon

- Typically we just deal with the ball $B_{\leq 1}(0)$.
- For a metric valued field K , $\{x \in K : |x| \leq 1\}$ is a subring - the *valuation ring*.
- This is a metric structure in the ring language.

Metric Valued Fields, à Rideau-Kikuchi, Scanlon, Simon

- Typically we just deal with the ball $B_{\leq 1}(0)$.
- For a metric valued field K , $\{x \in K : |x| \leq 1\}$ is a subring - the *valuation ring*.
- This is a metric structure in the ring language.
- Unfortunately, these are elementarily equivalent to other convex subrings of metric valued fields.

Ordered Real Closed Metric Valuation Rings

Theorem (A., Bejarano)

- *Models of our theory ORCMVR are convex subrings of real closed metric valued fields.*
- *These have quantifier-elimination once we add a divisibility relation.*
- *These can only be weakly o-minimal.*

Real Closed Metric Valued Fields à Ben Yaacov

- Itai Ben Yaacov studied the projective line instead

Real Closed Metric Valued Fields à Ben Yaacov

- Itai Ben Yaacov studied the projective line instead
- We need to view these as **cyclic** orders

Real Closed Metric Valued Fields à Ben Yaacov

- Itai Ben Yaacov studied the projective line instead
- We need to view these as **cyclic** orders

Theorem (A., Bejarano)

Any model of ORCMVF (the projective line) is an o-minimal expansion of a model of UDCO, the complete theory of ultrametric dense cyclic orders.

Further Directions

- (Weakly) ω -minimal expansions of MLOs are **distal**, and thus admit **distal cell decompositions**.

Further Directions

- (Weakly) o-minimal expansions of MLOs are **distal**, and thus admit **distal cell decompositions**.
Can we strengthen this to o-minimal cell decomposition?

Further Directions

- (Weakly) ω -minimal expansions of MLOs are **distal**, and thus admit **distal cell decompositions**.
Can we strengthen this to ω -minimal cell decomposition?
- If $U_S \models \text{UDLO}$ was \aleph_0 -categorical, its extremely amenable automorphism group would mean its age is Ramsey.

Further Directions

- (Weakly) ω -minimal expansions of MLOs are **distal**, and thus admit **distal cell decompositions**.
Can we strengthen this to ω -minimal cell decomposition?
- If $U_S \models \text{UDLO}$ was \aleph_0 -categorical, its extremely amenable automorphism group would mean its age is Ramsey.
What can we say for an **approximately** \aleph_0 -categorical structure?

Thank you, Rutgers!